

Healthcare system resilience and adaptability to pandemic disruptions in the United States

Received: 4 December 2023

Accepted: 31 May 2024

Published online: 02 July 2024

 Check for updatesLu Zhong^{1,2,4}, Dimitri Lopez^{1,4}, Sen Pei³ & Jianxi Gao^{1,2}✉

Understanding healthcare system resilience has become paramount, particularly in the wake of the COVID-19 pandemic, which imposed unprecedented burdens on healthcare services and severely impacted public health. Resilience is defined as the system's ability to absorb, recover from and adapt to disruptions; however, despite extensive studies on this subject, we still lack empirical evidence and mathematical tools to quantify its adaptability (the ability of the system to adjust to and learn from disruptions). By analyzing millions of patients' electronic medical records across US states, we find that the COVID-19 pandemic caused two successive waves of disruptions within the healthcare systems, enabling natural experiment analysis of the adaptive capacity of each system to adapt to past disruptions. We generalized the quantification framework and found that the US healthcare systems exhibit substantial adaptability ($\rho = 0.58$) but only a moderate level of resilience ($r = 0.70$). When considering system responses across racial groups, Black and Hispanic groups were more severely impacted by pandemic disruptions than white and Asian groups. Physician abundance was the key characteristic for determining healthcare system resilience. Our results offer vital guidance in designing resilient and sustainable healthcare systems to prepare for future waves of disruptions akin to COVID-19 pandemics.

Global crises such as climate change, environmental pollution, conflicts or global pandemics continue to pose great challenges to the healthcare system^{1,2}. These challenges are not solely a result of the escalating scale of these crises but also their prolonged duration with successive disruptions. The COVID-19 pandemic is a convergence of these dual challenges—an unprecedentedly large-scale crisis that has persisted for over 2 years, occurred in multiple waves and resulted in millions of hospitalizations and over 1.2 million deaths in the United States (US; Fig. 1a). The pandemic has led to disruptions to routine medical services, stemming from factors such as public fear of infections during visits to healthcare facilities³, stay-at-home policies⁴, patient access to care and

limited supply of services. This has led to delays and cancellations in non-COVID-19 emergency services and essential care⁴, and the inability to maintain essential services led to adverse and lasting consequences. For instance, in the US, 9.4 million cancer screenings, and treatments were either delayed or canceled due to the pandemic⁵ and the maternal mortality rate increased from 0.017% in 2019 to 0.032% deaths in 2021 (refs. 6, 7). Furthermore, the pandemic has disproportionately impacted marginalized groups such as people of color, low-income populations and those with underlying health conditions^{8,9}.

To improve overall public health outcomes and mitigate the negative consequences, it is important to enhance the resilience of

¹Department of Computer Science, Rensselaer Polytechnic Institute, Troy, NY, USA. ²Network Science and Technology Center, Rensselaer Polytechnic Institute, Troy, NY, USA. ³Department of Environmental Health Sciences, Columbia University, New York City, NY, USA. ⁴These authors contributed equally: Lu Zhong, Dimitri Lopez. ✉e-mail: jianxi.gao@gmail.com




Fig. 1 | Adaptive responses to successive disruptions in healthcare systems.

a, As seen in the data on COVID-19 hospitalizations, the pandemic brought successive disruptions to healthcare systems. **b**, The adaptive response cycle within the healthcare system involves multiple cycles of disruption and recovery, leading to enhanced resilience through learning from previous disruptions. By examining the system's performance in maintaining non-COVID-19 patient visits (**c–f**), we assess its adaptability and resilience during successive disruptions. With two disruptions as the example in **c**, when the disruption event i (for example, onset of the COVID-19 pandemic and second wave of the outbreak) occurs at time $t_{s,i}$

the healthcare system's performance (non-COVID-19 visits) begins to decline, reaches a negative peak and then returns to the target performance level at $t_{r,i}$. Each disruption i undergoes a disruption and recovery phase, characterized by disruption amplitude (α_i), disruption duration ($T_i = t_{r,i} - t_{s,i}$), disruption rate (u_i) and recovery rate (v_i). **c**, A system with low resilience and low adaptability. **d**, A system with high resilience is characterized by smaller amplitude/duration. **e**, A system with high adaptability is characterized by its ability to better absorb disruptions $u_{i+1} < u_i$ or quicker recover $v_{i+1} > v_i$ during the second disruption. **f**, A system with high resilience and high adaptability.

the healthcare system to maintain essential health services despite disruptions^{10–12}. Substantial research has been conducted in this area, including investigations into the conceptual framework of healthcare systems^{13–19}, examination of the impact of COVID-19 pandemic on disrupting health service delivery and degrading healthcare quality^{2,4,20–22} and proposals of measures, such as the social vulnerability index (SVI)^{23,24} and preparedness index (PPI)^{25,26}. However, so far, no standard definition and measurement for healthcare system resilience have been established²⁷.

Resilience is typically defined as the system's ability to absorb and recover from every single disruption^{28–31} and also the system's ability to adapt to multiple successive disruptions^{27,32} (Supplementary Table 1). Existing research on system resilience, spanning various fields including healthcare, ecology, business and industry, predominantly focuses on the static aspect of absorbing and recovery that enables a system to bounce back after the disruption^{33–36}. However, its ability to adapt to disruptions from disasters is rarely explored. As illustrated in Fig. 1b, mentioned in refs. 32,37, adaptability is defined as an adaptive cycle marked by recurrent disruptions and the innate system capability to learn from previous disturbances, ultimately leading to a more resilient system³⁸. As recurrent disruptions are rarely observed and recorded, the practical observation and quantitative assessment of this adaptive cycle in real-world systems remain absent. The COVID-19 pandemic, which results in successive waves of disruptions, provides a natural experiment to study the adaptability in a real-world healthcare system.

In this study, we analyzed millions of patients' records using electronic medical record (EMR) data (Methods; Supplementary Tables 2 and 3) to create a comprehensive assessment of healthcare systems' resilience and adaptability to disruptions caused by the COVID-19 pandemic in the US. We examined 23 essential healthcare services over a broad range of health needs (Methods; Supplementary Table 4), including chronic disease care (for example, Alzheimer's disease, cancer and heart diseases) and maternal care (for example, pregnancy). For the healthcare system in each state, we first identified the number of disruptions that the healthcare system encountered and quantified their durations and amplitudes. By comparing a system's performance in absorbing and recovering across disruptions, we were then able to evaluate the system's adaptability and compare it across various health services and distinct patient groups. We also quantified the total system resilience by assessing the loss of patient visits due to the pandemic. Using the COVID-19 pandemic as an example, this study offered a quantification framework for assessing healthcare resilience and adaptability. Table 1 summarizes the findings and policy implications.

Results

Adaptive responses to successive disruptions in healthcare systems

The resilience of healthcare systems measures the collective response of diverse entities, such as healthcare providers, hospitals, insurers, pharmacists, the general public and government entities, in sustaining the uninterrupted provision of essential services for patients^{4,18},

Table 1 | Policy summary

Background	Resilience is defined as the system's ability to absorb and recover from each single disruption, as well as its adaptability—to adapt or transform itself to better respond to multiple successive disruptions. However, despite the COVID-19 pandemic bringing multiple waves of disruptions to healthcare systems, there has been limited research dedicated to quantifying their adaptability to successive disruptions and their resilience during the pandemic.
Main findings and limitations	By analyzing extensive EMR data across US states, we find that the COVID-19 pandemic led to two successive disruptions within healthcare systems. We generalized the quantification framework and assessed the resilience of healthcare systems across various states for different essential services and populations according to race and ethnicity. The results show that healthcare systems demonstrate substantial adaptability but only a moderate level of resilience. Services for chronic disease treatment exhibit higher resilience compared to maternal services. Black and Hispanic populations were most affected by severe disruptions when compared to white and Asian groups. By examining the relationship between system resilience and factors such as pandemic severity, physician shortages and socioeconomic variables, we identified physician abundance as the pivotal characteristic influencing healthcare system responses. Limitations of this study include bias in the data collection process and missing attributes within the EMR dataset, potentially compromising the accuracy and reliability of predictive trends related to patient visits based on historical data.
Policy implications	Our results highlight the importance of improving the system's adaptability to effectively respond to ongoing disruptions, especially for maternal care, minority populations in the US and states with a scarcity of physicians, high poverty and low employment. The resilience and adaptability indexes we have introduced, founded on a dynamic perspective, complement existing metrics like the SVI and PPI, offering guidance for future disasters that spike waves of disruptions akin to COVID-19 pandemics.

as visualized in Fig. 1b. As the COVID-19 pandemic has progressed, it has given rise to several new virus variants, some of which have triggered multiple disruptions in health services. However, the existing quantification framework has primarily focused on a single disruption³⁵. Here we extend the quantification framework to multiple disruptions. Specifically, if there are no external disruptions, the system will maintain its expected performance $P(t)$, represented by non-COVID-19 patient visits for essential services. As illustrated in the example in Fig. 1c, the actual performance features two disruptions. The dynamics of every single disruption i can be captured by its disruption amplitude α_i (disruption severity), duration T_i , disruption rate u_i and recovery rate v_i (equation (1) and Supplementary Fig. 1). Disruption amplitude and duration gauge the severity of a disruption; higher values mean a more severe impact. Meanwhile, the disruption rate reflects the system's management of the disruption's progress, and the recovery rate indicates how efficiently the system returns to normalcy. Using Fig. 1c as a comparison, a more resilient healthcare system (Fig. 1d) can minimize amplitude and duration, ultimately leading to a smaller loss of patient visits. A higher adaptive healthcare system (Fig. 1e) can slow current disruption rates (or increase recovery rates) than previous ones, as evidenced by $u_2 < u_1$ or $v_2 > v_1$. In Fig. 1f, a healthcare system exhibits both high adaptability and resilience. Please refer to the Methods for further detailed measurements of the resilience index (r) and adaptability index (ρ).

Figure 2a shows the trend of patient visits to essential services in US states. During the prepandemic period (2017–2020), patient visits increased steadily. This increase can be attributed to the increased adoption of certified EMR technology by US hospitals, coupled with the Affordable Care Act's expansion of healthcare resources. We used a

predictive model, which incorporates the real-world increasing adoption of EMR technology among physicians within the data, to estimate the expected number of patient visits if the COVID-19 pandemic had not occurred (Methods; Extended Data Fig. 1 and Supplementary Figs. 2–5). In comparison to the expected patient visits $P(t)$, the observed patient visits $O(t)$ started to fluctuate and then decrease sharply at around the beginning of 2020 in Fig. 2a. Among the 49 analyzed states, 40 of them encountered two consecutive disruptions (Extended Data Table 1). The initial disruption generally occurred between January 2020 and May 2021, followed by a second disruption from June 2021 to the end of 2022. The two disruptions are highly correlated to the waves of the COVID-19 pandemic, with the initial disruption corresponding to the onset of the pandemic, and the second one being exacerbated by the emergence of new, more contagious variants (such as the Omicron variant in the last quarter of 2021). The disruption duration aligns with the results obtained from external datasets on emergency department visits and hospital discharges (Extended Data Fig. 2). All these findings of two disruptions contrast with previous studies that assumed that the COVID-19 pandemic caused a single disruption^{2,4,22}.

To quantify the system's adaptive response, it first needs to decode each disruption. Figures 2 and 3 illustrate the characteristics of the two disruptions. On average, the second disruption tends to have a longer duration and a larger amplitude than the initial disruption. However, the disruption rate during the second event was lower than that of the first event. The comparison between these two disruptions suggests that the healthcare system primed during the initial disruption can absorb disruption to decelerate its disruption rates. Regarding recovery, 25 states do not return to expected levels during the first disruption, while 38 of them remain unrecovered until the end of 2022 during the second disruption (Extended Data Table 1). As the recovery rate is largely unknown, we use the disruption rate to measure the system's adaptability. As depicted in Fig. 2b, systems with a disruption rate smaller than that of the first disruption tend to exhibit higher adaptability. Systems experiencing greater amplitude and extended durations in both disruptions tend to have lower resilience.

Figure 2c,d displays the ranking of adaptability and resilience of healthcare systems in US states. Most states exhibit a positive adaptability index, indicating their ability to improve during the second disruption. Five states stood out with notably negative adaptability indices, suggesting that states didn't have enough resources to better prepare for the following disruptions. Michigan achieves the highest resilience scores with nearly no loss of patient visits ($r = 0.98$), while Wyoming and Louisiana have the lowest resilience scores ($r = 0.48$ and $r = 0.41$). States that exhibit high adaptivity tend to also demonstrate high resilience (Pearson coefficient = 0.24, P value around 0.09; Supplementary Fig. 6). This reflects states with high adaptability being better equipped to handle subsequent disruptions, ultimately resulting in less loss of patient visits during the whole period and high resilience. When comparing the indices with the SVI^{23,24}, we find that the resilience indices are negatively correlated with the SVI for the state (Pearson coefficient = -0.38 , P value around 0.001; Extended Data Fig. 3). These results provide a reasonable validity of the resilience indices. States characterized by higher social vulnerability to disasters tend to have lower resilience scores.

Adaptability and resilience among essential services

To explore healthcare system resilience and adaptability in more detail, we group patient visit data based on the services that patients received for their disease. Among essential health services, we focused on chronic disease care and maternal care within the availability of our dataset. For the states with records exceeding a threshold of 1,000 patients in both of these services, 82.6% of chronic disease care and 100% of maternal care experienced two disruptions (Extended Data Table 1). For chronic disease care, 52.2% of states did not achieve recovery by the end of the first disruption, and 83.6% remained




Fig. 2 | Adaptability and resilience assessment of US healthcare systems.

a, Temporal trend of non-COVID-19 patient visits across US states, from January 2017 to December 2022. Among the 49 analyzed states, 40 (81.63%) experienced two successive disruptions. The initial disruption generally occurred between January 2020 and May 2021, followed by a second disruption starting in June 2021, and 38 of them were not recovered by the end of 2022. **b**, Disruption amplitude a_i , duration T_i and disruption rate u_i characterize each disruption for

states in high or low levels of adaptability and resilience. **c**, State rankings for adaptability with index ρ in the range of $(-1, 1)$, where a positive value indicates a capacity to adapt. **d**, State rankings for resilience with index $r \in (0, 1)$, where a higher value indicates a greater capacity to encounter disruptions and sustain the volume of visits. For better comparison, the plots in **c** and **d** are in descending order of resilience index. A system is considered to have high adaptability when $\rho > 0.5$ and high resilience when $r > 0.7$, highlighted in a gray band.

unrecovered in the second disruption, which extends until the end of 2022. For maternal care, a more severe situation was observed, with 66.6% of states failing to recover during the initial disruption and 86.8% of states still not recovering in the second disruption.

As illustrated in Fig. 3, during the second disruption, health-care services demonstrated substantial disruption amplitude and

prolonged durations. Their enhanced capacity to mitigate disruptions in the second phase results in an overall positive adaptability index for the healthcare system across US states ($\rho = 0.58 \pm 0.05$). Maternal care displayed less adaptability ($\rho = 0.48 \pm 0.10$), in contrast to chronic disease care, which displayed higher adaptability ($\rho = 0.61 \pm 0.06$). For overall resilience, the healthcare system across US states exhibited a




Fig. 3 | Adaptability and resilience assessment for essential services.

To measure healthcare system responses for essential services, we categorize non-COVID-19 patient visits into the following two specific services across US states: chronic disease care and maternal care. The ‘all’ category encompasses all services. **a**, The average adaptability (ρ). **b**, The average resilience (r).

c, d, Parameters characterize each disruption for services. The parameters include **(c)** disruption rate (u_i) and recovery rate (v_i) and **(d)** disruption amplitude (α_i) and disruption duration (T_i). The adaptability and resilience indices and parameters across all states are expressed by the average, along with the lower and upper limits of the 95% CI. CI, confidence level.

moderate level of resilience ($r = 0.70 \pm 0.03$). Maternal care thus exhibited lower resilience ($r = 0.74 \pm 0.03$) and experienced disruptions of greater duration, while chronic disease treatment exhibited higher resilience ($r = 0.76 \pm 0.03$). Supplementary Fig. 4 shows the sensitivity analysis of predictive models across various healthcare services. Supplementary Fig. 7 shows the resilience and adaptability indexes of subservices like chronic obstructive pulmonary disease (COPD), cancer, heart disease and diabetes. Extended Data Fig. 4 shows the results for services for dialysis.

Adaptability and resilience among patient groups

Besides maintaining essential services, a resilient healthcare system should be able to provide adequate care to all patients, regardless of their socioeconomic status, race or ethnicity. We performed subgroup analyses according to race and ethnicity, to measure the number of their visits during the COVID-19 pandemic. More than 22 states experienced two disruptions across all patient groups (Extended Data Table 1).

Using data from New York State as an example (Fig. 4a), across all patient groups, there is a decrease in disruption rates during the second wave of disruption, demonstrating positive adaptability. The Asian population had the lowest adaptability index ($\rho_{\text{Asian}} = 0.36 \pm 0.10$), while the white population had the highest adaptability index ($\rho_{\text{white}} = 0.56 \pm 0.06$; see Fig. 4b). This suggests that the Asian population is well-prepared to mitigate the impact of the first disruption, indicating limited room for further improvement and adaptability. The Asian population experienced a minimal decline in patient visits and the shortest duration, highlighting the highest resilience score. Meanwhile, the Hispanic and Black populations encountered severe disruptions in amplitude and duration, exhibiting the lowest resilience index. When considering all states (Fig. 4c–e), the Asian population garners the highest resilience score ($r_{\text{Asian}} = 0.80 \pm 0.02$), followed by the white population ($r_{\text{white}} = 0.74 \pm 0.03$). Conversely, the Black and Hispanic populations present the least amount of resilience with scores of $r_{\text{Black}} = 0.73 \pm 0.03$ and $r_{\text{Hispanic}} = 0.72 \pm 0.03$, respectively. Supplementary Fig. 4 shows the sensitivity analyses of predictive models across populations according to race.

Association with pandemic severity, physician shortages and socioeconomic factors

The different performance of the healthcare system in US states during the COVID-19 pandemic is complex and multifactorial, with a range of factors, including pandemic severity, healthcare infrastructure, resources, lockdown policies, socioeconomic factors and the political climate all having a role. To incorporate these factors into healthcare system planning and decision-making processes, we need to sort out their effects on the resilience index r , adaptability index ρ and disruption amplitude α . We thus extensively collect state-level COVID-19 cases, physician abundance and socioeconomic factors to estimate their correlations (Methods). As presented in Table 2, there was a positive correlation between states’ resilience index and the abundance of physicians (0.313 ($P = 0.001$)), while there were negative correlations between resilience index and local poverty levels (-0.328 ($P = 0.001$)) and also the unemployment rates (-0.156 ($P = 0.001$)). We also observed significant correlations between the adaptability index and these factors. For the two pandemic disruptions, their amplitudes exhibit negative associations with physician abundance (-0.241 ($P = 0.005$); -0.205 ($P = 0.005$)) and positive associations with local poverty levels (0.253 ($P = 0.003$); 0.234 ($P = 0.015$)) and uninsured levels (0.333 ($P = 0.001$); 0.401 ($P = 0.001$)). The results indicate that states with low physician abundance, high poverty rates, high unemployment and low insurance coverage are at a higher risk of severe disruption during the pandemic, presenting smaller healthcare system resilience and adaptability.

Discussion

By analyzing EMR data across US states, we measure the number of patient visits to essential services to analyze the collective response of diverse entities within the healthcare system in the face of consecutive disruptions during the COVID-19 pandemic. Our quantification framework of resilience encompasses key metrics such as the resilience index, adaptability index and parameters that describe the amplitude and duration of each disruption, as well as the system performance in managing disruption and recovery rate. Our findings reveal that the healthcare system underwent two waves of disruptions, demonstrating an




Fig. 4 | Adaptability and resilience assessment for patient groups by race and ethnicity. To measure healthcare system responses for patients by race and ethnicity, we group non-COVID-19 visits according to patients' attributes across US states. **a**, Illustration of the temporal trend of non-COVID-19 patient visits for Asian, Black, Hispanic and white groups in the New York state. **b**, The average adaptability (ρ). **c**, The average resilience (r) of patient groups. **d,e**, The

parameters characterize each disruption. The parameters include (**d**) disruption rate (u_i) and recovery rate (v_i) and (**e**) disruption amplitude (α_i) and disruption duration (T_i). The adaptability and resilience indices and parameters across all states are expressed by the average, along with the lower and upper limits of the 95% confidence level.

adaptive response wherein lessons from the initial disruption enhance the system's capacity to absorb disruptions and expedite recovery. Over 90% of the healthcare systems performed better during the second disruption. As of the end of 2022, about 77% of healthcare systems have yet to fully rebound to normal levels. Consistent with previous research⁴, state demographic attributes, such as high poverty levels and high rates of unemployment, were correlated with low resilience and adaptability. The abundance of the physician workforce has an

essential role in determining healthcare resilience and adaptability^{39–41}. The findings highlight the importance of strategically organizing the physician workforce during disasters and enhancing collaborations across states⁴².

By examining different healthcare services, we found that chronic disease care is more resilient and adaptive than maternal care. This is because chronic disease care needs long-term treatments and is more flexible in delivery across primary care clinics, specialty clinics and

Table 2 | Pearson correlation coefficients assessing the relationships between system adaptability/resilience and pandemic severity, physician shortages and socioeconomic factors in US states

Parameters	COVID-19 cases	Physician per 100,000	Poverty percentile	Unemployment percentile	Uninsurance percentile	Age ≥65 percentile	Age ≤17 percentile	Minority percentile
Adaptability index	−0.033 (<i>P</i> =0.507)	0.102 (<i>P</i> = 0.039)	− 0.082 (<i>P</i> = 0.046)	− 0.146 (<i>P</i> = 0.003)	−0.010 (<i>P</i> =0.828)	0.040 (<i>P</i> =0.404)	−0.009 (<i>P</i> =0.847)	−0.0046 (<i>P</i> =0.350)
Resilience index	−0.054 (<i>P</i> =0.278)	0.313 (<i>P</i> = 0.001)	− 0.328 (<i>P</i> = 0.001)	− 0.156 (<i>P</i> = 0.001)	− 0.47 (<i>P</i> = 0.001)	0.116 (<i>P</i> = 0.019)	− 0.209 (<i>P</i> = 0.001)	− 0.217 (<i>P</i> = 0.009)
Amplitude α (first disruption)	0.114 (<i>P</i> =0.198)	− 0.241 (<i>P</i> = 0.005)	0.253 (<i>P</i> = 0.003)	0.093 (<i>P</i> =0.295)	0.333 (<i>P</i> = 0.001)	−0.116 (<i>P</i> =0.191)	0.175 (<i>P</i> = 0.047)	0.054 (<i>P</i> =0.369)
Amplitude α (second disruption)	0.132 (<i>P</i> =0.178)	− 0.205 (<i>P</i> = 0.035)	0.234 (<i>P</i> = 0.015)	0.206 (<i>P</i> = 0.034)	0.401 (<i>P</i> = 0.001)	−0.042 (<i>P</i> =0.668)	0.143 (<i>P</i> =0.144)	0.103 (<i>P</i> =0.292)

Significant correlations, determined by a two-sided test and indicated by a *P* value less than the threshold of 0.05, are given in bold.

home health services, adding resilience through diverse ways of care⁴. Meanwhile, maternal care relies on short-term specialized services, its acute nature makes it more vulnerable to external disruptions⁴³. Among the subservices for chronic disease care (Supplementary Fig. 7), services for heart disease management demonstrate lower resilience and adaptability compared to services for conditions such as asthma, COPD, cancer and diabetes. These findings emphasize the pressing requirement for heightened assistance and focus on heart disease care during a pandemic.

We also examined the healthcare resilience of populations by race and ethnicity. We found that the pandemic severely affected Black and Hispanic populations, leading to harsher disruptions and lower resilience indexes for these populations. These disparities are likely rooted in socioeconomic inequalities that Black and Hispanic communities typically encounter greater obstacles in accessing healthcare services, particularly during external disruptions^{44,45}. The findings underscore the importance of enhancing chronic disease care and maternal care to alleviate the exacerbation of inequality, especially for Black and Hispanic communities during crises.

During the COVID-19 pandemic, the number of patient visits to healthcare providers was jointly affected by the supply of services and patient demand. The availability of resources can limit access to healthcare services, and patient's perception of infection risk may reduce the demand for nonessential services. Our analysis considered the impact of both the supply- and demand-side factors. Using essential service dialysis as an illustration, we disentangle the two factors and offer a rigorous assessment of the availability of delivering services from healthcare infrastructure and resources (the supply side). Patients undergoing dialysis usually need to attend several sessions each week, and their treatment is dependent on consistent and uninterrupted access to these services. Patient visits to dialysis services exhibit two distinct disruptions, differing in duration from other healthcare services. The first disruption occurred from December 2019 to March 2022, while the second disruption commenced in April 2022 with no observed recovery. The dialysis service exhibits delayed disruptions and higher resilience ($r = 0.89$). The results suggest that the healthcare system still underwent two disruptions without substantial change in service demand, indicating that the finding of two successive disruptions is replicated.

Nevertheless, the quantification framework has several limitations. First, although the EMR dataset we used is extensive within the US, the incomplete collection and biased data sampling in the dataset can affect the accuracy of our assessment. Specifically, the dataset may primarily represent states that have broadly adopted and implemented the EMR technology, potentially introducing bias into our analysis, particularly in regions where EMR adoption was low. Second, the missing attribute in the dataset can also potentially introduce bias into the analysis. Efforts to mitigate biases in data collection methodologies^{46,47} and integrating additional data sources can further validate

the findings. Third, our assessment relies on the appropriate selection of a predictive model for forecasting expected patient visits. While we conduct various sensitivity analyses using alternative models, a more sophisticated model is needed to account for uncertainty in data and state variations within the system. Addressing structural uncertainty by considering multiple potential models and assessing their implications for predictions will also be needed for more comprehensive and reliable assessment.

Several avenues merit exploration for future research for a more comprehensive quantification of healthcare system resilience. It is crucial to consider excessive recovery as a new dimension of system resilience, aiming not only to return to its original state but to recover lost visits and progress toward a more optimal state (Supplementary Fig. 8). It is necessary to untangle the intertwined impacts of successive disruptions, particularly when disruptions stem from different crises. Furthermore, there is also a necessity to broaden the scope beyond COVID-19 waves to comprehend healthcare system performance across different crises, especially for future guiding healthcare system resilience against more deadly diseases associated with climate change. For example, conducting a comparative analysis of responses to the 2003 severe acute respiratory syndrome (SARS) outbreak and the 2019 COVID-19 pandemic could serve as a guide. Additionally, while our measure reflects the overall resilience of the healthcare system, it is crucial to exclude demand factors to concentrate solely on the dynamics of service availability when evaluating the resilience for specific hospitals and other healthcare facilities. Moreover, our macroscopic measures are limited in offering higher-resolution guidance on each system component. Future studies should delve into sophisticated computational methods and consider additional factors such as the intricate connections between healthcare facilities, resource management, the healthcare workforce, supply chain dynamics and the quality of patient care^{18,41}. Finally, our measures can only be assessed retrospectively. Future endeavors should strive to provide real-time estimates for proactive decision-making.

In summary, our study provides a quantification framework for healthcare system resilience that can be applied generically to various healthcare services, patient groups and different regions^{48,49}. The proposed adaptability index and resilience index allow the characterization of healthcare system performance during and across multiple disruptions. These two indices can also offer a valuable complement to existing crisis management tools, such as SVI²⁴ and PPI^{25,26}, which predominantly focus on either the socioeconomic status of a region or the availability of healthcare resources before disruption strikes. In brief, this framework could provide policymakers with the essential insights to make informed adaptations to successive disruptions caused by prolonged disasters^{17,50,51}.

Online content

Any methods, additional references, Nature Portfolio reporting summaries, source data, extended data, supplementary information,

acknowledgements, peer review information; details of author contributions and competing interests; and statements of data and code availability are available at <https://doi.org/10.1038/s41591-024-03103-6>.

References

- World Health Organization. Maintaining essential health services: operational guidance for the COVID-19 context. www.who.int/publications/i/item/WHO-2019-nCoV-essential_health_services-2020.2 (2020).
- Doubova, S. V., Leslie, H. H., Kruk, M. E., Pérez-Cuevas, R. & Arsenault, C. Disruption in essential health services in Mexico during COVID-19: an interrupted time series analysis of health information system data. *BMJ Glob. Health* **6**, e006204 (2021).
- World Health Organization. Pulse survey on continuity of essential health services during the COVID-19 pandemic. www.who.int/publications/i/item/WHO-2019-nCoV-EHS_continuity-survey-2020.1 (2020).
- Arsenault, C. et al. COVID-19 and resilience of healthcare systems in ten countries. *Nat. Med.* **28**, 1314–1324 (2022).
- Chen, R. C., Haynes, K., Du, S., Barron, J. & Katz, A. J. Association of cancer screening deficit in the United States with the COVID-19 pandemic. *JAMA Oncol.* **7**, 878–884 (2021).
- Hoyert, D. L. Maternal mortality rates in the United States, 2022. NCHS Health E-Stats. www.cdc.gov/nchs/data/hestat/maternal-mortality/2022/maternal-mortality-rates-2022.pdf (2024).
- Chmielewska, B. et al. Effects of the COVID-19 pandemic on maternal and perinatal outcomes: a systematic review and meta-analysis. *Lancet Glob. Health* **9**, e759–e772 (2021).
- Chowkwanyun, M. & Reed, A. L. Jr. Racial health disparities and COVID-19—caution and context. *N. Engl. J. Med.* **383**, 201–203 (2020).
- Lopez, L., Hart, L. H. & Katz, M. H. Racial and ethnic health disparities related to COVID-19. *JAMA* **325**, 719–720 (2021).
- Balabanova, D. et al. Good health at low cost 25 years on: lessons for the future of health systems strengthening. *Lancet* **381**, 2118–2133 (2013).
- Kruk, M. E. et al. High-quality health systems in the sustainable development goals era: time for a revolution. *Lancet Glob. Health* **6**, e1196–e1252 (2018).
- Budd, J. et al. Digital technologies in the public-health response to COVID-19. *Nat. Med.* **26**, 1183–1192 (2020).
- Wulff, K., Donato, D. & Lurie, N. What is health resilience and how can we build it? *Annu. Rev. Public Health* **36**, 361–374 (2015).
- Aitsi-Selmi, A., Egawa, S., Sasaki, H., Wannous, C. & Murray, V. The Sendai framework for disaster risk reduction: renewing the global commitment to people's resilience, health, and well-being. *Int. J. Disaster Risk Sci.* **6**, 164–176 (2015).
- Mirzoev, T. & Kane, S. What is health systems responsiveness? Review of existing knowledge and proposed conceptual framework. *BMJ Glob. Health* **2**, e000486 (2017).
- Turenne, C. P. et al. Conceptual analysis of health systems resilience: a scoping review. *Soc. Sci. Med.* **232**, 168–180 (2019).
- Biddle, L., Wahedi, K. & Bozorgmehr, K. Health system resilience: a literature review of empirical research. *Health Policy Plan.* **35**, 1084–1109 (2020).
- Haldane, V. et al. Health systems resilience in managing the COVID-19 pandemic: lessons from 28 countries. *Nat. Med.* **27**, 964–980 (2021).
- Iflaifel, M., Lim, R. H., Ryan, K. & Crowley, C. Resilient health care: a systematic review of conceptualisations, study methods and factors that develop resilience. *BMC Health Serv. Res.* **20**, 324 (2020).
- Roy, C. M. et al. Assessing the indirect effects of COVID-19 on healthcare delivery, utilization and health outcomes: a scoping review. *Eur. J. Public Health* **31**, 634–640 (2021).
- Doubova, S. V. et al. Overcoming disruptions in essential health services during the COVID-19 pandemic in Mexico. *BMJ Glob. Health* **7**, e008099 (2022).
- Kiarie, H. et al. The COVID-19 pandemic and disruptions to essential health services in Kenya: a retrospective time-series analysis. *Lancet Glob. Health* **10**, e1257–e1267 (2022).
- Flanagan, B. E., Gregory, E. W., Hallisey, E. J., Heitgerd, J. L. & Lewis, B. A social vulnerability index for disaster management. *J. Homel. Secur. Emerg. Manag.* **8**, (2011).
- Centers for Disease Control and Prevention. CDC social vulnerability index. www.arcgis.com/apps/InteractiveLegend/index.html?appid=4d92395405c844a49a36eea90aa6bbfb (2018).
- Zhang, F. et al. Predictors of COVID-19 epidemics in countries of the World Health Organization African Region. *Nat. Med.* **27**, 2041–2047 (2021).
- Chaudhry, R., Dranitsaris, G., Mubashir, T., Bartoszko, J. & Riazi, S. A country level analysis measuring the impact of government actions, country preparedness and socioeconomic factors on COVID-19 mortality and related health outcomes. *EClinicalMedicine* **25**, 100464 (2020).
- Kruk, M. E. et al. Building resilient health systems: a proposal for a resilience index. *BMJ* **357**, j2323 (2017).
- Fisher, L. More than 70 ways to show resilience. *Nature* **518**, 35–35 (2015).
- Linkov, I. et al. Changing the resilience paradigm. *Nat. Clim. Change* **4**, 407–409 (2014).
- Linkov, I. & Trump, B. D. *The Science and Practice of Resilience* (Springer, 2019).
- Comfort, L. K., Boin, A. & Demchak, C. C. *Designing Resilience: Preparing for Extreme Events* (University of Pittsburgh Press, 2010).
- Holling, C. S., & Gunderson, L. H. Resilience and adaptive cycles. In *Panarchy: Understanding Transformations in Human and Natural Systems* Ch. 2 (2002).
- Liu, X. et al. Network resilience. *Phys. Rep.* **971**, 1–108 (2022).
- Poulin, C. & Kane, M. B. Infrastructure resilience curves: performance measures and summary metrics. *Reliab. Eng. Syst. Saf.* **216**, 107926 (2021).
- Cassottana, B., Aydin, N. Y. & Tang, L. C. Quantitative assessment of system response during disruptions: an application to water distribution systems. *J. Water Resour. Plan. Manag.* **147**, 04021002 (2021).
- Gao, J., Barzel, B. & Barabási, A.-L. Universal resilience patterns in complex networks. *Nature* **530**, 307–312 (2016).
- Holling, C. S. Resilience and stability of ecological systems. *Annu. Rev. Ecol. Syst.* **4**, 1–23 (1973).
- Zhang, H., Liu, X., Wang, Q., Zhang, W. & Gao, J. Co-adaptation enhances the resilience of mutualistic networks. *J. R. Soc. Interface* **17**, 20200236 (2020).
- Association of American Medical Colleges. The complexities of physician supply and demand: projections from 2019 to 2034. digirepo.nlm.nih.gov/master/borndig/9918417887306676/9918417887306676.pdf (2019).
- Sturmberg, J. & Lanham, H. J. Understanding health care delivery as a complex system: achieving best possible health outcomes for individuals and communities by focusing on interdependencies. *J. Eval. Clin. Pract.* **20**, 1005–1009 (2014).
- Lo Sardo, D. R. et al. Quantification of the resilience of primary care networks by stress testing the health care system. *Proc. Natl Acad. Sci. USA* **116**, 23930–23935 (2019).
- Achour, N. & Price, A. D. Resilience strategies of healthcare facilities: present and future. *Int. J. Disaster Resil. Built. Environ.* **1**, 264–276 (2010).

43. U.S. Government Accountability Office. Maternal health: outcomes worsened and disparities persisted during the pandemic. www.gao.gov/products/gao-23-105871 (2022).
 44. Abedi, V. et al. Racial, economic, and health inequality and COVID-19 infection in the United States. *J. Racial Ethn. Health Disparities* **8**, 732–742 (2021).
 45. Bamba, C., Riordan, R., Ford, J. & Matthews, F. The COVID-19 pandemic and health inequalities. *J. Epidemiol. Community Health* **74**, 964–968 (2020).
 46. Aubinet, M., Vesala, T. & Papale, D. (eds). *Eddy Covariance: A Practical Guide To Measurement and Data Analysis* (Springer, 2012).
 47. Dahlen, A. & Charu, V. Analysis of sampling bias in large health care claims databases. *JAMA Netw. Open* **6**, e2249804 (2023).
 48. Mills, A. Health care systems in low-and middle-income countries. *N. Engl. J. Med.* **370**, 552–557 (2014).
 49. Yabe, T., Rao, P. S. C., Ukkusuri, S. V. & Cutter, S. L. Toward data-driven, dynamical complex systems approaches to disaster resilience. *Proc. Natl Acad. Sci. USA* **119**, e2111997119 (2022).
 50. Ceferino, L., Mitrani-Reiser, J., Kiremidjian, A., Deierlein, G. & Bambarén, C. Effective plans for hospital system response to earthquake emergencies. *Nat. Commun.* **11**, 4325 (2020).
 51. Saulnier, D. D. et al. A health systems resilience research agenda: moving from concept to practice. *BMJ Glob. Health* **6**, e006779 (2021).
- Publisher's note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.
- Springer Nature or its licensor (e.g. a society or other partner) holds exclusive rights to this article under a publishing agreement with the author(s) or other rightsholder(s); author self-archiving of the accepted manuscript version of this article is solely governed by the terms of such publishing agreement and applicable law.
- © The Author(s), under exclusive licence to Springer Nature America, Inc. 2024

Methods

Data

The EMR dataset we used, referred to as the Healthjump dataset, is provided by the COVID-19 Research Database⁵². Healthjump is a data integration platform that exports EMR and practice management systems. It aggregates data from over 70,000 hospitals and clinics, as well as more than 1,500 healthcare organizations, covering all states in the US. With records dating back to 1995, the dataset encompasses information on over 33 million patients and 1 million healthcare providers. It includes a wide array of patient data, including diagnoses, procedures, encounters and medical histories, sourced from participating members of the Healthjump network. Additionally, it contains patients' sociodemographic attributes, such as race, sex and others. The dataset details, statistics and limitations are presented in Supplementary Tables 2 and 3.

In examining the correlation between states' resilience/adaptability index with COVID-19 infections, physician abundance and sociodemographic factors, we first gather cumulative infection cases in each state from the Johns Hopkins Coronavirus Resource Center⁵³. We collect physician abundance data regarding physician numbers for each state, drawing from the 2019 State Physician Workforce Data⁵⁴. The sociodemographic factors analyzed include poverty levels, unemployment rates, uninsured levels, the proportion of the youth (less than 17), the proportion of the elderly (greater than 65) and the proportion of minority populations, all sourced from the CDC/ATSDR SVI²⁴.

Ethics statement

Ethical approval was not required for this study, as the data used for analysis was from the fully anonymized Healthjump EMR database⁵². The database complies with the Health Insurance Portability and Accountability Act of 1996, ensuring the protection of patient information through strict privacy policies and agreements with patients and healthcare providers. Therefore, ethical approval was not needed, as the database has no identifiable information about individual patients.

Demographic information

Demographic information was restricted to race and ethnicity. For the EMR-reported race and ethnicity, patients were designated by themselves or by a healthcare provider. For race, patients were designated as 'American Indian or Alaska Native', 'Asian', 'Black or African American', 'white', 'Native Hawaiian or other Pacific Islander' and 'other race'. For ethnicity, patients were designated as 'Hispanic or Latino', 'not Hispanic or Latino' and 'unknown'. For simplification, we consider the main race/ethnic categories in the study.

Essential health services of concern

Essential health services encompass vital healthcare provisions crucial for enhancing and preserving public health^{1,4}. These services typically comprise maternal care, chronic disease management, diagnostic and laboratory services, vaccination, primary care, emergency care and others. Due to limitations in data availability and the exclusion of data related to care for patients with COVID-19, our study focuses solely on chronic disease care and maternal care across a spectrum of 23 diseases (Supplementary Table 4). We analyzed the absolute number of visits for these services. Additionally, we normalized the data to mitigate the impact of state population size and policies related to joining Healthjump's EMR system in our analyses.

Quantification framework for successive disruptions

We used mathematical models that provide key parameters to characterize the system's behavior during disruption and recovery processes. While numerous models exist⁵⁵, the beta family equations uniquely offer flexibility (Supplementary Fig. 1). Building upon the framework that describes the system's behavior under a single disruption³⁵, we generalize this framework to multiple successive disruptions, as given below:

$$O(t) = \begin{cases} P(t); & 0 \leq t < t_{s_1} \\ P(t) - \alpha_1 \frac{(\theta_1 + \vartheta_1)^{\theta_1 + \vartheta_1}}{\theta_1^{\theta_1} + \vartheta_1^{\vartheta_1}} \left(\frac{t}{T_1} \right)^{\theta_1} \left(1 - \frac{t}{T_1} \right)^{\vartheta_1}; & t_{s_1} \leq t < t_{r_1} \\ P(t) - \alpha_2 \frac{(\theta_2 + \vartheta_2)^{\theta_2 + \vartheta_2}}{\theta_2^{\theta_2} + \vartheta_2^{\vartheta_2}} \left(\frac{t}{T_2} \right)^{\theta_2} \left(1 - \frac{t}{T_2} \right)^{\vartheta_2}; & t_{s_2} \leq t < t_{r_2} \\ \dots & \dots \\ P(t); & t > t_{r_n} \end{cases}, \quad (1)$$

where $O(t)$ is the actual observed performance of patient visits and $P(t)$ is the predicted performance if the pandemic didn't occur. Suppose there are n disruptions. Each disruption i has an amplitude α_i , which is the scale factor, defined as the severity of disruption on the system. The other two parameters θ_i and ϑ_i determine the curve's shape for disruption and recovery respectively, as reflected in the system's ability to manage the processes. Within the same duration T_i ($T_i = t_{r_i} - t_{s_i}$), if $\theta_i = \vartheta_i$, a symmetric disruption and recovery occur; if $\theta_i > \vartheta_i$, a slow disruption is followed by a fast recovery; if $\theta_i < \vartheta_i$, a fast disruption is followed by a slow recovery. We define disruption rate as $u_i = \frac{1}{\theta_i T_i}$ and its recovery rate as $v_i = \frac{1}{\vartheta_i T_i}$. A smaller value of θ_i or ϑ_i leads to quick disruption or recover. A shorter duration T_i results in a rapid deterioration of performance to its maximum extent.

To assess the system's performance across n disruptions, we introduce the adaptability index in terms of disruption rate across consecutive disruptions i and $i + 1$,

$$\rho = \frac{1}{n} \sum_{i=1}^n \frac{-(u_{i+1} - u_i)}{\max(u_{i+1}, u_i)}, \quad (2)$$

where $\rho \in (-1, 1)$. Specifically, $\rho > 0$ indicates that the system exhibits adaptability with $u_{i+1} < u_i$, signifying that the rate of disruption $i + 1$ is smaller than that of disruption i . Conversely, $\rho < 0$ suggests that the system lacks adaptability with $u_{i+1} > u_i$, indicating that the rate of disruption $i + 1$ is larger than that of disruption i . Higher values of ρ signify an increased level of adaptability of the system. In the case of a single disruption, $\rho = 1$. The adaptability index can also be measured in terms of recovery rate. As the recovery rate is largely unknown in our results, we only use the disruption rate to measure the system adaptivity.

Following the classic way^{29,30,55}, we measure system resilience in terms of performance loss as

$$r = 1 - \frac{\int_{t_s}^{t_r} (P(t) - O(t)) dt}{\int_{t_s}^{t_r} P(t) dt}, \quad (3)$$

where $r \in (0, 1)$. The term $\int_{t_s}^{t_r} (P(t) - O(t)) dt$ is the total loss between expected $P(t)$ and observed patient visits $O(t)$. The division by integral of expected performance $\int_{t_s}^{t_r} P(t) dt$ normalizes the results to a range between 0 and 1. A value of 1 indicates no performance loss (perfect resilience), while 0 indicates a complete loss.

Predictive model for quantification framework

For the analysis of healthcare resilience during the COVID-19 pandemic, we extracted data from 2017 to the end of 2022. Commencing from January 2020, the onset of the pandemic, we designate this as the starting date, denoted as t_s . The dataset is subsequently divided into the following two distinct periods: the prepandemic period ($t < t_s$, from January 2017 to December 2019) and the pandemic period ($t \geq t_s$, from January 2020). With the growing adoption of EMR technology in US hospitals and increased accessibility to healthcare facilities due to the Affordable Care Act, there has been a noticeable rise in patient visit volumes in our datasets. To capture the expected patient visits in equation (1), we leverage the number of physicians in the data that participate

in EMR technology, state population and physician-to-population ratio³⁴ to assess expected patient visits,

$$P(t) = \mathcal{P}_0(t)\sigma n, \quad (4)$$

where $\mathcal{P}_0(t)$ is the monthly number of physicians that adopted EMR technology in the state, n is the state population and σ is the state physician-to-population ratio, adjusted by monthly visit frequency in the prepandemic period. Through the models, we can predict the expected patient visits that digital health platforms would accumulate if no disruptions occurred beyond t_s . We also use the innovation (EMR) adoption model⁵⁶ and time-series model⁵⁷, that is, the generalized logistic model and exponential smoothing model, and consider the seasonality for the sensitivity test. The comparisons of predictive models are provided in Extended Data Fig. 1 and Supplementary Figs. 2–5. To smooth out seasonal fluctuations and identify underlying trends or patterns, we use the 3-month moving average on data. For comparison across states and services, the volume of patient visits at $t = 0$ is normalized to 1.

Identification of disruptions

To fit the observed performance $O(t)$ with equation (1), we first identify the number of disruptions n and the duration T_i of each disruption i . The disruptions are identified through segmented least squares, which enables the detection of both the peak of decline and subsequent increase in performance and divides the duration T_i into disruption and recovery phases. Disruptions failing to meet the following criteria will be excluded: (1) those lasting less than 3 months (where $T_i < 3$) and (2) those with a minimum performance loss ratio below 5%. These two criteria are used to exclude disruptions with not significant losses and those attributed to data fluctuations caused by seasonal effects (see Supplementary Fig. 5 for the sensitivity test of criteria). Then we infer the parameters α_i , θ_i and ϑ_i for the observed performance for each disruption i by using the iterative optimization method^{58,59}.

Statistical analysis

To demonstrate the resilience and adaptability of various services and races, we represent their average across analyzed states with the lower and upper limits of 95% confidence level in Figs. 3 and 4. For a total of M analyzed state, the average of resilience for services is calculated as $\bar{r}^i = \frac{\sum_m r_m^i}{M}$, where r_m^i represents the resilience at state m for services i . Similarly, the average of resilience for racial groups is calculated as $\bar{r}^j = \frac{\sum_m r_m^j}{M}$, where r_m^j represents the resilience at state m for race j . The same calculations are applied for the average of adaptability.

We use the Pearson correlation coefficient to gauge the correlation between state resilience (or adaptability) with various factors, such as COVID-19 infections, physician abundance, sociodemographic factors and SVI. This coefficient, ranging between -1 and 1 , quantifies both the strength and direction of the relationship between two variables.

Reporting summary

Further information on research design is available in the Nature Portfolio Reporting Summary linked to this article.

Data availability

The EMR dataset that supports the findings of this study is available from the Healthjump database, provided by the COVID-19 Research Database Consortium (<https://covid19researchdatabase.org/>). However, restrictions apply to accessing these data, which were used under license for the current study. The EMR dataset is not publicly available. The data on COVID-19 infection cases in each state are collected from the Johns Hopkins Coronavirus Resource Center (<https://github.com/CSSEGISandData/COVID-19>). The general physician abundance data regarding physician numbers in each state are collected from the 2019

State Physician Workforce Data (<https://www.aamc.org/data-reports/workforce/report/state-physician-workforce-data-report>). The sociodemographic factors in each state are collected from the CDC/ATSDR SVI (<https://www.atsdr.cdc.gov/placeandhealth/svi/index.html>). For validation, external summary datasets on patient visits to physicians, emergency departments and the number of hospital discharges during the pandemic are sourced from the National Center for Health Statistics (<https://www.cdc.gov/nchs/index.htm>) and the US Census Bureau (<https://www.census.gov/>). For the results dashboard, please see the website ResilienceHealthSys.com.

Code availability

The code used in the study for the quantification framework is available at https://github.com/lucinezhong/healthcare_resilience_quantification_framework.

References

52. The COVID-19 Research Database. covid19researchdatabase.org/ (accessed 1 January 2023).
53. Dong, E., Du, H. & Gardner, L. An interactive web-based dashboard to track COVID-19 in real time. *Lancet Infect. Dis.* **20**, 533–534 (2020).
54. Association of American Medical Colleges. State physician workforce data report. www.aamc.org/data-reports/workforce/report/state-physician-workforce-data-report (2019).
55. Cassottana, B., Shen, L. & Tang, L. C. Modeling the recovery process: a key dimension of resilience. *Reliab. Eng. Syst. Saf.* **190**, 106528 (2019).
56. Rogers, E. M., Singhal, A. & Quinlan, M. M. *An Integrated Approach to Communication Theory and Research* (Routledge, 2014).
57. Hamilton, J. D. *Time Series Analysis* (Princeton University Press, 2020).
58. Kelley, C. T. *Iterative Methods for Optimization* (SIAM, 1999).
59. Strutz, T. *Data Fitting and Uncertainty: A Practical Introduction to Weighted Least Squares and Beyond* (Springer, 2011).

Acknowledgements

The data, technology and services used in generating these research findings were generously supplied pro bono by the COVID-19 Research Database partners, who are acknowledged at <https://covid19researchdatabase.org/>. This work was supported, in whole or in part, by the Bill & Melinda Gates Foundation (No. CORONAVIRUSHUB-D-21-00120). Under the grant conditions of the Foundation, a Creative Commons Attribution 4.0 Generic License has already been assigned to the Author Accepted Manuscript version that might arise from this submission. We also acknowledge the support of Research Accelerator grants funded by the USA National Science Foundation (2047488), the Rensselaer-IBM AI Research Collaboration, the USA National Science Foundation (DMS-2229605) and Centers for Disease Control and Prevention (U01CK000592).

Author contributions

L.Z., D.L. and J.G. conceived the project and designed the study. D.L. and L.Z. performed the data analyses. L.Z. and D.L. wrote the first draft of the manuscript. L.Z., S.P. and J.G. contributed to interpreting the results and improving the manuscript. J.G. was the lead writer of the manuscript.

Competing interests

The authors declare no competing interests.

Additional information

Extended data is available for this paper at <https://doi.org/10.1038/s41591-024-03103-6>.


Supplementary information The online version contains supplementary material available at <https://doi.org/10.1038/s41591-024-03103-6>.

Correspondence and requests for materials should be addressed to Jianxi Gao.

Peer review information *Nature Medicine* thanks Catherine Arsenault and the other, anonymous, reviewer(s) for their contribution to

the peer review of this work. Primary Handling Editor: Ming Yang, in collaboration with the *Nature Medicine* team.


Reprints and permissions information is available at www.nature.com/reprints.



National	Model-1	Model-2	Model-3	Model-4	Model-0
First disruption period	202001-202105	202001-202105	202001-202105	202001-202105	202001-202105
Second disruption period	202107-202208	202106-202209	202106-unknown	202106-unknown	202106-unknown
Resilience	0.79	0.63	0.44	0.61	0.73
Adaptability	0.27	0.30	0.53	0.43	0.42


Extended Data Fig. 1 | Comparison of predictive models $P(t)$ using national volume of Non-COVID-19 patient visits in Healthjump dataset. Comparison of predictive models $P(t)$ using national volume of Non-COVID-19 patient visits of Healthjump dataset. (a) Estimation model incorporating physician EMR technology adoption counts within Healthjump database. (b) Generalized

logistic model utilizing AHA (American Hospital Association) EMR adoption rates. (c) Simple exponential smoothing model integrated with AHA EMR adoption rates. (d) Seasonal exponential smoothing model incorporating AHA EMR adoption rates. (e) Constant population who have active visits during the pre-pandemic period. All models identify two disruptions.




Extended Data Fig. 2 | Volume of non-COVID-19 patient care in the US from 2019 to 2022 provided by other datasets. (a) Patient visits to physicians, sourced from the National Center for Health Statistics. **(b)** Patient visits to emergency departments, also sourced from the National Center for Health Statistics. **(c)** Number of hospital discharges, provided by the US Census Bureau.

Compared against a baseline represented by a blue line, indicative of average patient visits before the pandemic, the observed patient visits (represented by a red line) show two disruptions, marked by dashed vertical lines. The first disruption spans from the first quarter of 2020 to the second/third quarter of 2021, while the second disruption begins in the third/fourth quarter of 2021.



Extended Data Fig. 3 | Resilience index of states versus social vulnerability index. (a) State resilience index. (b) State social vulnerability index. (c) Pearson correlation coefficient. The resilience index is negatively correlated with the social vulnerability index. The correlation significance is determined by a two-sided test and indicated by a *P* value.



Extended Data Fig. 4 | Temporal trend of patient visits for dialysis service. Similar to conclusions drawn for other essential services, two disruptions are observed, but the period differs, with the first occurring from December 2019 to March 2022 and the second beginning in April 2022 with no observed recovery.

Extended Data Table 1 | Successive disruptions on healthcare system from 2020 to 2022

Percentage	Disruptions		Not recovered (1st disruption)	Not recovered (2nd disruption)	States
	Once	Twice			
All	18.3%	81.63%	51.0%	77.5%	49
Chronic Disease Care	17.3%	82.6%	52.2%	82.6%	23
Maternal Care	0%	100%	66.6%	86.8%	15
Asian Population	31.2%	68.7%	37.5%	56.2%	32
Black Population	18.7%	81.2%	53.1%	78.1%	32
Hispanic Population	17.1%	83.8%	60.0%	77.1%	35
White Population	25%	74%	47.2%	65.9%	44

Successive disruptions on US healthcare systems from 2020 to 2022. We classify the healthcare system as ‘not recovered’ if the observed non-COVID-19 patient visits keep less than 95% of the expected counts.

Reporting Summary

Nature Portfolio wishes to improve the reproducibility of the work that we publish. This form provides structure for consistency and transparency in reporting. For further information on Nature Portfolio policies, see our [Editorial Policies](#) and the [Editorial Policy Checklist](#).

Statistics

For all statistical analyses, confirm that the following items are present in the figure legend, table legend, main text, or Methods section.

n/a	Confirmed
<input checked="" type="checkbox"/>	<input type="checkbox"/> The exact sample size (n) for each experimental group/condition, given as a discrete number and unit of measurement
<input checked="" type="checkbox"/>	<input type="checkbox"/> A statement on whether measurements were taken from distinct samples or whether the same sample was measured repeatedly
<input checked="" type="checkbox"/>	<input type="checkbox"/> The statistical test(s) used AND whether they are one- or two-sided <i>Only common tests should be described solely by name; describe more complex techniques in the Methods section.</i>
<input checked="" type="checkbox"/>	<input type="checkbox"/> A description of all covariates tested
<input type="checkbox"/>	<input checked="" type="checkbox"/> A description of any assumptions or corrections, such as tests of normality and adjustment for multiple comparisons
<input type="checkbox"/>	<input checked="" type="checkbox"/> A full description of the statistical parameters including central tendency (e.g. means) or other basic estimates (e.g. regression coefficient) AND variation (e.g. standard deviation) or associated estimates of uncertainty (e.g. confidence intervals)
<input type="checkbox"/>	<input checked="" type="checkbox"/> For null hypothesis testing, the test statistic (e.g. F , t , r) with confidence intervals, effect sizes, degrees of freedom and P value noted <i>Give P values as exact values whenever suitable.</i>
<input checked="" type="checkbox"/>	<input type="checkbox"/> For Bayesian analysis, information on the choice of priors and Markov chain Monte Carlo settings
<input checked="" type="checkbox"/>	<input type="checkbox"/> For hierarchical and complex designs, identification of the appropriate level for tests and full reporting of outcomes
<input type="checkbox"/>	<input checked="" type="checkbox"/> Estimates of effect sizes (e.g. Cohen's d , Pearson's r), indicating how they were calculated

Our web collection on [statistics for biologists](#) contains articles on many of the points above.

Software and code

Policy information about [availability of computer code](#)

Data collection	No software was used for data collection.
Data analysis	No commercial and open-source codes are used. Our team develops the Python code for analyzing the volume of patient visits and measuring system resilience and adaptivity. The code is provided at https://github.com/lucinezhong/healthcare_resilience_quantification_framework .

For manuscripts utilizing custom algorithms or software that are central to the research but not yet described in published literature, software must be made available to editors and reviewers. We strongly encourage code deposition in a community repository (e.g. GitHub). See the Nature Portfolio [guidelines for submitting code & software](#) for further information.

Data

Policy information about [availability of data](#)

All manuscripts must include a [data availability statement](#). This statement should provide the following information, where applicable:

- Accession codes, unique identifiers, or web links for publicly available datasets
- A description of any restrictions on data availability
- For clinical datasets or third party data, please ensure that the statement adheres to our [policy](#)

The electronic medical record dataset that supports the findings of this study is available from the Healthjump database, provided by the COVID-19 Research Database consortium (<https://covid19researchdatabase.org/>). However, restrictions apply to the availability of these data, which were used under license for the current study. The EMR dataset is not publicly available.

The data on COVID-19 infection cases in each state is collected from the Johns Hopkins Coronavirus Resource Center (<https://github.com/CSSEGISandData/COVID-19>).

The general physician abundance data regarding physician numbers in each state is collected from the 2019 State Physician Workforce Data (<https://www.aamc.org/data-reports/workforce/report/state-physician-workforce-data-report>).

The social factors include poverty levels, unemployment rates, uninsured levels, the proportion of the youth (less than 17), the proportion of the elderly (greater than 65), and the proportion of minority populations are collected from the CDC/ATSDR Social Vulnerability Index (<https://www.atsdr.cdc.gov/placeandhealth/svi/index.html>).

For validation, external summary datasets on patient visits to physicians, emergency departments, and the number of hospital discharges during the pandemic are supplied by the National Center for Health Statistics (<https://www.cdc.gov/nchs/index.htm>) and the US Census Bureau (<https://www.census.gov/>).

Research involving human participants, their data, or biological material

Policy information about studies with [human participants or human data](#). See also policy information about [sex, gender \(identity/presentation\), and sexual orientation](#) and [race, ethnicity and racism](#).

Reporting on sex and gender

Use the terms sex (biological attribute) and gender (shaped by social and cultural circumstances) carefully in order to avoid confusing both terms. Indicate if findings apply to only one sex or gender; describe whether sex and gender were considered in study design; whether sex and/or gender was determined based on self-reporting or assigned and methods used.

Provide in the source data disaggregated sex and gender data, where this information has been collected, and if consent has been obtained for sharing of individual-level data; provide overall numbers in this Reporting Summary. Please state if this information has not been collected.

Report sex- and gender-based analyses where performed, justify reasons for lack of sex- and gender-based analysis.

Reporting on race, ethnicity, or other socially relevant groupings

Please specify the socially constructed or socially relevant categorization variable(s) used in your manuscript and explain why they were used. Please note that such variables should not be used as proxies for other socially constructed/relevant variables (for example, race or ethnicity should not be used as a proxy for socioeconomic status).

Provide clear definitions of the relevant terms used, how they were provided (by the participants/respondents, the researchers, or third parties), and the method(s) used to classify people into the different categories (e.g. self-report, census or administrative data, social media data, etc.)

Please provide details about how you controlled for confounding variables in your analyses.

Population characteristics

Describe the covariate-relevant population characteristics of the human research participants (e.g. age, genotypic information, past and current diagnosis and treatment categories). If you filled out the behavioural & social sciences study design questions and have nothing to add here, write "See above."

Recruitment

Describe how participants were recruited. Outline any potential self-selection bias or other biases that may be present and how these are likely to impact results.

Ethics oversight

Identify the organization(s) that approved the study protocol.

Note that full information on the approval of the study protocol must also be provided in the manuscript.

Field-specific reporting

Please select the one below that is the best fit for your research. If you are not sure, read the appropriate sections before making your selection.

☐ Life sciences ☒ Behavioural & social sciences ☐ Ecological, evolutionary & environmental sciences

For a reference copy of the document with all sections, see nature.com/documents/nr-reporting-summary-flat.pdf

Behavioural & social sciences study design

All studies must disclose on these points even when the disclosure is negative.

Study description

This study is a quantitative analysis that harnesses billions of electronic health records to assess the resilience and adaptability of healthcare systems during the COVID-19 pandemic. This quantitative framework aims to identify disruption timelines and evaluate system performance in delivering essential services such as chronic disease treatment and maternal care across various racial and ethnic population groups.

Research sample

The dataset utilized in this study comprises over one billion electronic medical records spanning the period from 2017 to 2022 across 50 US states. Among these records, approximately 30 millions contain ICD codes related to services for chronic illness treatment and maternal care. The breakdown of records by racial and ethnic groups is as follows: one million (0.5%) for Asian, twenty million (10.4%) for Black, twenty million (10.4%) for Hispanic, and one hundred and fifty million (78%) for White patients. The dataset also encompasses additional demographic attributes, such as age. It comprises three age groups: youth (0-19 years, 9%), middle-aged (19-60 years, 53%), and elderly individuals (60-100 years, 38%).

To compare the resilience and adaptivity index across states, along with their respective social factors such as poverty levels, unemployment rates, uninsured levels, proportions of youth (under 17), elderly (over 65), and minority populations, we gathered data from the CDC/ATSDR Social Vulnerability Index (SVI) available at <https://www.atsdr.cdc.gov/placeandhealth/svi/index.html>.

Sampling strategy

No sampling strategy is used for analysis.

Data collection	We collect monthly number of patient visits , their demographic information (e.g., race and ethnicity) , and the ICD-10 code in their diagnosis at each state from the Healthjump database. The Healthjump database collection relies on comprehensive integration of facilities that join EMR technology. Concerning personal demographic information, the database is committed to complying with all pertinent federal and state laws and adheres strictly to the Privacy Policy. It establishes agreements with healthcare providers to ensure the transfer of patient information to the database is in accordance with HIPAA (Health Insurance Portability and Accountability Act of 1996) regulations.
Timing	The data we studies is monthly data from 2017/01 to 2022/12.
Data exclusions	NONE
Non-participation	NONE
Randomization	Not relevant in this study. This study measures outcomes without any manipulation or intervention by researchers. When exploring the correlation between resilience and adaptivity index with social factors in each state, randomization is irrelevant.

Reporting for specific materials, systems and methods

We require information from authors about some types of materials, experimental systems and methods used in many studies. Here, indicate whether each material, system or method listed is relevant to your study. If you are not sure if a list item applies to your research, read the appropriate section before selecting a response.

Materials & experimental systems

n/a	Involved in the study
<input checked="" type="checkbox"/>	<input type="checkbox"/> Antibodies
<input checked="" type="checkbox"/>	<input type="checkbox"/> Eukaryotic cell lines
<input checked="" type="checkbox"/>	<input type="checkbox"/> Palaeontology and archaeology
<input checked="" type="checkbox"/>	<input type="checkbox"/> Animals and other organisms
<input checked="" type="checkbox"/>	<input type="checkbox"/> Clinical data
<input checked="" type="checkbox"/>	<input type="checkbox"/> Dual use research of concern
<input checked="" type="checkbox"/>	<input type="checkbox"/> Plants

Methods

n/a	Involved in the study
<input checked="" type="checkbox"/>	<input type="checkbox"/> ChIP-seq
<input checked="" type="checkbox"/>	<input type="checkbox"/> Flow cytometry
<input checked="" type="checkbox"/>	<input type="checkbox"/> MRI-based neuroimaging

Plants

Seed stocks	Report on the source of all seed stocks or other plant material used. If applicable, state the seed stock centre and catalogue number. If plant specimens were collected from the field, describe the collection location, date and sampling procedures.
Novel plant genotypes	Describe the methods by which all novel plant genotypes were produced. This includes those generated by transgenic approaches, gene editing, chemical/radiation-based mutagenesis and hybridization. For transgenic lines, describe the transformation method, the number of independent lines analyzed and the generation upon which experiments were performed. For gene-edited lines, describe the editor used, the endogenous sequence targeted for editing, the targeting guide RNA sequence (if applicable) and how the editor was applied.
Authentication	Describe any authentication procedures for each seed stock used or novel genotype generated. Describe any experiments used to assess the effect of a mutation and, where applicable, how potential secondary effects (e.g. second site T-DNA insertions, mosaicism, off-target gene editing) were examined.